Technical Information Bulletin

BASF
The Chemical Company

Headline AMP® Fungicide

Northwest Kansas On-Farm Trials



Average of 18 trials (2011-2012) located in Hoxie, St Francis, Scott City and Brewster, Kansas.

Replicated University Studies



Average of 12 studies (2011-2013). **Locations:** Fort Collins, CO; Bethune, CO; Scottsbluff, NE; Lingle, WY; Farmington, NM. **Universities:** Colorado State University, University of Nebraska-Lincoln, University of Wyoming, New Mexico State University.

Best Use Recommendations

- Timing: VT (Full tassel) to R3 (Milk)
- Use Rate: 10 fl oz/A
- Carrier Volume: at least 2 GPA by air
- PHI: 20 days for field corn and 7 days for silage
- Adjuvants:
  - No adjuvants from V8 to VT
  - VT to R3-COC @ 1 pint/A or HSOC at 0.5 pint/A
  - No adjuvants if tank mixing with foliar fertilizer
- Foliar fertilizer:
  - No more than 50% of the total carrier volume
  - e.g. if total carrier volume is 2 GPA, then you should not use more than 1 gal/A of foliar fertilizer
- Miticide/Insecticide:
  - Headline AMP fungicide may be mixed with labeled miticides and insecticides.
  - Follow the most restrictive adjuvant recommendations for the tank-mix partners.

Headline AMP®
Fungicide

Always read and follow label directions.
Headline AMP is a registered trademark of BASF.
©2014 BASF Corporation. All Rights Reserved. APN# 1407010 Headline AMP-Corn-KS

BASF
The Chemical Company