

Pristine®

Fungicide

Managing Strawberry Botrytis Grey Mold in Florida

Benefits of Pristine® Fungicide

- Pristine fungicide offers excellent control of Powdery Mildew and Anthracnose when used at the 18.5 to 23 oz/Acre rates
- In Florida, Botrytis strains with reduced sensitivity to Pristine have been observed and the distribution of these strains varies across the Florida strawberry growing region
- Botrytis control with Pristine may be reduced if insensitive isolates are present

Time Period	December/January	February/March
Target Diseases	Anthracnose, Powdery Mildew; Susceptible Botrytis Grey Mold	Anthracnose, Powdery Mildew; Susceptible Botrytis Grey Mold
Pristine Use Recommendation	Apply at 18.5 to 23 oz/A	Apply at 23 oz/A as a rotational partner for products with different chemistries

Pristine Use Recommendations

To limit the potential for resistance development, rotate Pristine with Botrytis fungicides with different modes of action (Captec®, Captevate®, Inspire Super™, Switch®, Elevate®, or Scala®) and restrict consecutive applications of Pristine.

Always follow FRAC guidelines when using fungicides.



Always read and follow label directions.

Pristine® is a registered trademark of BASF Corp.
Captec® 4L, Captevate® and Elevate® are registered trademarks of Arysta LifeScience North America, LLC.
Inspire Super™ is a trademark and Switch® is a registered trademark of the Syngenta Group Company.
Scala® is a registered trademark of Bayer.
©2012 BASF Corporation. All Rights Reserved. APN # 12-12-043