



The Standard For Residual Weed Control

Benefits of Zidua® SC Herbicide

Longer Lasting

- Up to two weeks longer residual weed control than other herbicides

Flexibility

- Low use rate: As little as 1/20 the use rate of other residual herbicides
- Labeled for celery, chickpeas, corn, cotton, onions, peanuts, potatoes, soybeans, sunflowers, wheat, and other crops

Broad Spectrum Grass & Broadleaf Weed Control

- Superior Palmer Amaranth and Waterhemp control
- Excellent Italian ryegrass and Poa (annual bluegrass) control



Zidua SC herbicide is a low rate formulation that provides long lasting residual weed control.

Waterhemp Species Weed Control Ratings*

Herbicide	Rating
Zidua Herbicide	Excellent
S-metolachlor	Fair to Good
Warrant® Herbicide	Fair to Good

*Soybean herbicide performance ratings, Winfield United 2019 Crop Protection Guide pp. 96-97

Palmer Amaranth Weed Control Ratings**

Herbicide	Rating
Zidua Herbicide	9
Dual II Magnum® Herbicide	8
Warrant Herbicide	8

**University of Arkansas, 2020. MP44 Recommended chemicals for weed and brush control, page 41.

















BASF

We create chemistry

Technical Information Bulletin

Zidua® SC Herbicide Timing and Use Rate by Crop

Crop	Timing	Use Rate (fl oz.)
 Celery†	1 to 6 days POST transplant, muck only	3.25
 Chickpea†	Preplant surface, PRE	2.5-3.25
 Corn	Preplant to V4	1.75-6.5
 Cotton	POST directed (layby) 5th leaf to beginning bloom	1.25-3.5
 Fallow	After harvest, before rotational crop interval	1.75-6.5
 Grass Seed†	Post-harvest to Jan 31, perennial ryegrass and tall fescue ¹	3.0
 Mint†	Dormant established stands in ID, MT, OR, and WA only	3.0
 Onion†	POST 2 to 6 leaves ²	2.0-2.75
 Peanuts	POST, at cracking to beginning pod	2.5-3.5
 Potato	PRE	2.5-3.25
 Safflower†	PRE	1.75-2.5
 Soybeans	Preplant to V6	1.75-5.75
 Sunflower	Preplant, PRE, POST 1 to 8 leaves	1.75-6.5
 Wheat	Delayed PRE to 4th Tiller ³	1.25-4.0

† Not labeled on coarse soils

¹ OR and WA only (see label for specific geography)

² Dry bulb onion and leek only, not green onions

³ PRE in ID, MT, OR, WA, 1.75-3.25 fl oz/A



To learn more about crop protection from BASF, visit www.agproducts.basf.us

BASF
We create chemistry

 **Zidua® SC**
Herbicide

Engenia Herbicide is a U.S. EPA Restricted Use Pesticide. Additional state restrictions may apply.

Always read and follow label directions. Zidua is a registered trademark of BASF. Dual II Magnum is a registered trademark of Syngenta. Warrant is a registered trademark of Monsanto Company. ©2021 BASF. All Rights Reserved.