



Bayer CropScience

**Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)**

PONCHO®/VOTiVO™

EPA Reg. No. 264-1109

For Use on Cotton

Supplemental Label

PONCHO®/VOTiVO™

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product package label for PONCHO®/VOTiVO™.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not apply where runoff is likely to occur. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on the label.

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

DIRECTIONS FOR USE

FOR USE AS A SEED TREATMENT

Apply PONCHO/VOTiVO as a commercial water-based slurry either alone or with other registered seed treatment insecticides and fungicides through standard slurry or mist commercial seed treatment equipment. Follow the most restrictive of the labeling limitations and precautions of all products used in mixtures. Treated seed must be labeled in accordance with the requirements of the Federal Seed Act.

APPLICATION RATES

CROP	INSECT PESTS	NEMATODE PESTS	RATE PONCHO/VOTiVO
Cotton, delinted, only	Cutworm, black	Columbia Lance Nematode Reniform Nematode Root Knot Nematode	2.4 fl oz / 100,000 seed (or 0.424 mg total ai / seed)

¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

RESTRICTIONS:

Regardless of type of application (seed treatment or foliar), do not apply more than 0.2lbs (Soybean), 0.1lbs (corn), or 0.2lbs (cotton) of active ingredient clothianidin per acre per season.

Resistance Management

The components in Poncho/VOTiVO belong to multiple chemistry classes. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Use labeled rates. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide/insecticide resistance management and/or integrated pest management recommendations for specific crops and pest populations. Bayer CropScience encourages responsible product stewardship to ensure effective long-term control of pests on this label.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with PONCHO/VOTiVO must be labeled with the following statements:

- This seed has been treated with PONCHO/VOTiVO, which contains 40.3% clothianidin **and 8.1% *Bacillus firmus* I-1582.**
- Do not use treated seed for food, feed, oil production or ethanol production.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with PONCHO/VOTiVO:

- Regardless of type of application (seed treatment or foliar), do not apply more than 0.2lbs (Soybean), 0.1lbs (corn), or 0.2lbs (cotton) of active ingredient clothianidin per acre per season.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound may be highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- For Cotton and Sweet corn: **Immediate Plant-back:** Cereal grains, Field corn, popcorn, sweet corn, cotton, rapeseed, soybeans, canola, root and tuber vegetables, **30-Day Plant-back:** Grasses, non-grass animal feeds, dried beans, **4-Month Plant-back:** Any crop without an earlier plant-back interval

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*