



Cevya® Fungicide on Carrot and Root Vegetables

Built to Last. Strong Control and Flexibility with Less Uncertainty.

Fast Uptake, High Potency and Residual Activity Leads to More Effective Disease Control.



Untreated




University of Florida
Standard Program




Cevya Fungicide
Program

2019-2020 University of Florida. Ben Broughton, Live Oak, FL. BASF sponsored trial and BASF images.

Effective *Alternaria* and *Cercospora* Control Even Under the Highest Pressure



Three trial average values GA (2) & CA (1) in 2017 (1) & 2018 (2) with adjuvants O-Si (1) or O-Si/MSO blend (2). Six trial average values adds GA (1), TX (1) & CA (1) in 2019 (3) with adjuvant O-Si/MSO blend (3). All trials BASF sponsored.



2016 TX, Adjuvant = NIS. 2017 TX, Adjuvant = O-Si. All trials BASF sponsored.



Cevya® Fungicide Best Use Recommendations

Application Tips

- Cevya fungicide should be applied preventively, prior to disease onset
- Thorough and uniform coverage for best performance
- Rainfast 1 hour after spray has dried

Pre-Harvest Interval: 7 days

Minimum Retreatment Interval: 7 days



Target Diseases	Use Rate (fl oz/A)	
	Single Application	Season Total
Alternaria leaf spot, (<i>Alternaria</i> spp.)	3-5	15
Cercospora leaf spot, (<i>Cercospora</i> spp.)		
Powdery mildew, (<i>Erysiphe</i> spp., <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.)		

Crop List – Root Vegetables* (Sub-Group 1B)

- Carrot

*For a full list of crops included in crop group, see label.



To learn more about crop protection products from BASF, visit www.agproducts.basf.us

BASF

We create chemistry

Cevya®
Fungicide