



Cevya® Fungicide on Cucurbit Vegetables


Built to Last. Strong Control and Flexibility with Less Uncertainty.

Fast Uptake, High Potency and Residual Activity Leads to More Effective Disease Control.



2018 Field Testing Sites in FL and GA. BASF Sponsored Research and BASF Images.

Effective Gummy Stem Blight Control Even Under the Highest Pressure



2019 GA. BASF Sponsored Trial. Gary Cloud GLC Consulting LLC. Tallahassee, FL. Watermelon var: Black Diamond. Four applications Oct 9, Oct 16, Oct 23 and Oct 29, 2019. All treatments included NIS adjuvant.



Cevya® Fungicide Best Use Recommendations

Application Tips

- Cevya fungicide should be applied preventively, prior to disease onset
- Thorough and uniform coverage for best performance
- Rainfast 1 hour after spray has dried

Pre-Harvest Interval: 0 days

Minimum Retreatment Interval: 7 days



Target Diseases	Use Rate (fl oz/A)	
	Single Application	Season Total
Gummy stem blight (<i>Didymella bryoniae</i>)	3-5	15
Powdery mildew (<i>Erysiphe</i> spp., <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.)		

Crop List – Cucurbit Vegetables* (Group 9)

- Cucumber; melon; pumpkin; squash

*For a full list of crops included in crop group, see label.



To learn more about crop protection products from BASF, visit www.agproducts.basf.us

BASF

We create chemistry

Cevya®
Fungicide