

Clubroot won't quit. Neither will we.

Clubroot is a constantly evolving disease in canola. As such, management practices need to evolve alongside the disease to minimize its impact today, as well as in the future. Sustainability of the US canola production industry guides our position on how to best manage clubroot and how we recommend deploying clubroot-resistant genetics. With all of this in mind, growers should implement an integrated pest management (IPM) strategy that includes:


- Canola **rotation** of a minimum of once every three years
- Taking **sanitation** steps to limit the movement of infected soil
- Controlling volunteer canola and other **brassica weeds** that can act as hosts for the disease
- Utilizing **patch management** to limit the movement of clubroot on your farm
- **Scouting** to identify the presence of the disease
- Clubroot-resistant **genetics** are only effective as a sustainable practice as part of an IPM strategy and should not be relied upon alone



Since their launch, InVigor® canola hybrids with clubroot resistance have been planted on over 51 million acres across Canada, where the clubroot disease pressure is much higher due to tighter rotations. That's more than any other seed company*. For 2025, US growers can choose from seven InVigor clubroot-resistant hybrids – all of which include our patented Pod Shatter Reduction technology.

InVigor L333PC, InVigor L350PC, InVigor Choice LR354PC, InVigor L340PC, InVigor L345PC and InVigor Choice LR344PC all contain the same **first-generation** clubroot resistance genetics. InVigor L343PC contains this resistance profile plus it contains **second-generation** clubroot resistance to additional emerging clubroot pathotypes to help defend against the ever-evolving pathogen.

To assist in determining the risk of clubroot on your farm and selecting the correct InVigor hybrid to grow, please refer to the following chart.



All agronomic recommendations include thorough scouting and implementing an integrated pest management strategy. ¹ When growing clubroot-resistant hybrids, we recommend using first-generation clubroot-resistant hybrids for two cycles OR until clubroot symptoms appear, whichever comes first, then consider switching to second-generation clubroot-resistant hybrids. All InVigor clubroot-resistant hybrids have been developed to be resistant to the most predominant clubroot pathotypes found in the US at the time of their registration.

*Source: 2019 and 2020 BPI (Business Planning Information) data

InVigor clubroot-resistant hybrids for 2025.

	DESCRIPTION	YIELD	TRAITS	STANDABILITY	MATURITY	GROWING ZONES	BLACKLEG
NEW InVigor L333PC	Introducing InVigor L333PC – our earliest maturing 300 series hybrid. With the InVigor performance you have come to trust, this hybrid is a great fit for those looking for an earlier-maturing hybrid. With high yield potential, good standability, our patented Pod Shatter Reduction technology and first-generation clubroot resistance ² , InVigor L333PC is a great fit for those in short and mid growing zones or if you are looking to beat the heat and get the combines rolling early. This hybrid demonstrated great performance in our 2023 Agronomic Excellence trials, making this an exciting new hybrid to try in 2025.	107.7% of the checks (InVigor L255PC and Pioneer [®] 45H33), BASF internal trials, 2022 & 2023, n=46 101.8% of InVigor L340PC, BASF internal trials, 2023, n=33 106.3% of InVigor L233P, BASF internal trials, 2023, n=33	Patented Pod Shatter Reduction technology First-generation clubroot resistance	Strong	½ day later than InVigor L233P	All growing areas in the US spring canola market	R Blackleg resistant
InVigor L350PC	Looking for a high-yielding hybrid in the mid to long growing areas? InVigor L350PC has it all. Our highest-yielding hybrid in our trials to date, this hybrid combines standability, first-generation clubroot resistance ² and our patented Pod Shatter Reduction technology.	108.8% of the checks (InVigor L255PC and Pioneer [®] 45H33), BASF internal trials, 2020 & 2021 108.9% of InVigor L340PC, 2021, n=22 114.3% of InVigor L233P, 2020 & 2021, n=36	Patented Pod Shatter Reduction technology First-generation clubroot resistance	Strong	4 ½ days later than InVigor L233P	All mid to long growing areas in the US spring canola market	R Blackleg resistant
InVigor ^{choice} LR354PC	InVigor Choice LR354PC offers growers high yield potential and great standability; with the added flexibility of harnessing both the LibertyLink [®] technology system and TruFlex [®] canola with Roundup Ready [®] Technology.	111.9% of the checks (InVigor L255PC and Pioneer [®] 45H33), BASF internal trials, 2020 & 2021 110.9% of InVigor Choice LR344PC, 2020 & 2021, n=33 112.5% of InVigor L233P, 2020 & 2021, n=33	Patented Pod Shatter Reduction technology First-generation clubroot resistance ² Both the LibertyLink technology system and TruFlex [®] canola with Roundup Ready [®] Technology	Strong	5 days later than InVigor L233P	All mid to long growing areas in the US spring canola market	R Blackleg resistant
InVigor L343PC	InVigor L343PC is a go-to choice when you want high yield potential in areas where you need second-generation clubroot resistance. ² Offers Pod Shatter Reduction and improved standability over InVigor L234PC. We recommend growing InVigor L343PC with second-generation clubroot resistance in clubroot-affected areas after two cycles of growing first-generation clubroot-resistant hybrids or when clubroot symptoms appear in first-generation clubroot-resistant hybrids.	111.3% of the checks (InVigor L233P and Pioneer [®]), WCC/RRC ³ trials, 2019 & 2020 106% of InVigor L233P, 2019 & 2020, n=43	Patented Pod Shatter Reduction technology Second-generation clubroot resistance	Strong	3 ½ days later than InVigor L233P	All growing areas in the US spring canola market	R Blackleg resistant
InVigor L340PC	InVigor L340PC has it all—high yield potential, Pod Shatter Reduction, first-generation clubroot resistance ² and strong standability. It is a great fit for growers who irrigate or when lodging is a concern.	108.9% of the new checks (InVigor L233P and Pioneer [®] 45H33), WCC/RRC ³ trials, 2019 107.8% of InVigor L233P trials, 2019, n=16	Patented Pod Shatter Reduction First-generation clubroot resistance	Strong	3 days later than InVigor L233P	All growing areas in the US spring canola market	R Blackleg resistant
InVigor L345PC	InVigor L345PC was the highest-yielding hybrid in our 2021 and 2022 Demonstration Strip Trials. Further improve the standability of InVigor L345PC by avoiding short rotations, fields with high volunteer canola populations, high levels of fertility or under irrigation.	111.9% of the checks (InVigor 5440 and Pioneer [®] 45H33), WCC/RRC ³ trials, 2017 & 2018 111.4% of InVigor L233P, 2018, n=28	Patented Pod Shatter Reduction technology First-generation clubroot resistance ²	Strong	3 ½ days later than InVigor L233P	All growing areas in the US spring canola market	R Blackleg resistant
InVigor ^{choice} LR344PC	InVigor Choice hybrid with Pod Shatter Reduction and first-generation clubroot resistance. ² InVigor Choice LR344PC ⁴ features both the LibertyLink technology system and TruFlex [®] canola with Roundup Ready [®] Technology. Perfect for growers looking to combine high yield potential InVigor canola genetics with the flexibility of Liberty [®] herbicide or Roundup [®] herbicide applications.	104.1% of the new checks (InVigor L233P and Pioneer [®] 45H33), WCC/RRC ³ trials, 2018 103.6% of InVigor L233P, 2018, n=12	Patented Pod Shatter Reduction technology First-generation clubroot resistance LibertyLink technology system and TruFlex [®] canola with Roundup Ready [®] Technology	Strong	2 ½ days later than InVigor L233P	All growing areas in the US spring canola market	R Blackleg resistant

Please note: Maturity and standability are based on performance ratings and data compiled from several InVigor internal trials over multiple years. Results may vary on your farm due to environmental factors and preferred management practices.

n=number of balanced trials.

²To predominant clubroot pathotypes found in Canada at the time of registration. InVigor L333PC, InVigor L350PC, InVigor Choice LR354PC, InVigor L340PC, InVigor Choice LR344PC and InVigor L345PC share the same first-generation clubroot resistance profile. InVigor L343PC has this resistance profile plus it contains second-generation multigenetic clubroot resistance to additional clubroot pathotypes to help combat evolving clubroot pathotypes. ³Western Canadian Canola/Rapeseed Recommending Committee. ⁴This product is approved for planting in Washington, Oregon, Idaho, Montana and the following counties of North Dakota: Divide, Williams, McKenzie. For additional information please contact your BASF representative.

Best management practices for clubroot.

While clubroot-resistant hybrids play an important role in managing the disease, they cannot be solely relied upon. It's important that we use them in combination with other strategies to prevent the breakdown of resistance and further spread of disease. That's why BASF continues to recommend an IPM strategy that complements the InVigor clubroot-resistant hybrids you're using on your farm.

Integrated pest management (IPM) with clubroot-resistant hybrids.

Sanitation.

Preventing spread of the disease is important for long-term control, even with clubroot-resistant hybrids. Sanitation practices are essential for fields known to be infested with clubroot; it's also important to ensure that areas without clubroot remain that way.

Keep spore loads low.

Extending canola rotations will help to prevent the buildup of clubroot resting spores. Paired with growing clubroot-resistant hybrids, you can reduce the potential for shifting populations to virulent pathotypes that could overcome resistance. In order to preserve existing clubroot resistance for as long as possible, a minimum two-year break from a host crop is required—and longer if inoculum levels are high.

Weed control.

Control of canola volunteers and other brassica weeds that act as hosts for clubroot will be essential to keeping the selection pressure low by not allowing the disease to build up in the non-canola years.

Scouting.

Even growers using clubroot-resistant hybrids should scout their field regularly to see if there is any evidence of the resistance breaking down. Also, if you're growing susceptible canola hybrids, scout for clubroot symptomology. If the field is known to be infested with clubroot, scouting should be thorough. In other words, you should examine several different areas of the field to see if there is any indication of resistance falling. If the field is not known to be infested, scouting near the field entry points should occur once near the end of the growing season. If small patches of clubroot are identified, it's crucial to take the time to hand pull and eliminate the galls since one heavily infected plant could produce up to 16 billion spores.

In general, all the prevention strategies should still be employed even with clubroot-resistant hybrids.



Frequently asked questions – clubroot in canola.

At BASF, our goal is to keep growers informed about the latest agronomic research information. That's why we're providing this question and answer section to address several key findings on managing clubroot in canola.

Q. Does growing an InVigor clubroot-resistant hybrid mean I will not see clubroot symptoms in my field?

A. Clubroot resistance protects the crop from being infected, but it does not control spores in the soil. That's why a low number of plants within the field may exhibit clubroot symptoms. These plants may be non-resistant volunteers from previous crops, off-types or potentially a new clubroot pathotype to which InVigor hybrids may be susceptible. For more information, reach out to your local BASF representative.

Q. What is the best time to grow second-generation clubroot-resistant hybrids?

A. Ideally, growers should start seeding a second-generation clubroot-resistant hybrid before symptoms appear in their first-generation hybrid. In fields where clubroot has been confirmed, second-generation hybrids should replace first-generation hybrids after two cycles (assuming at least a one-in-three-year rotation). Growers can transition to a second-generation hybrid earlier than this if symptoms appear in their first-generation hybrid.

Q. Will InVigor clubroot-resistant hybrids prevent the spread of clubroot to non-infested fields?

A. While growing InVigor clubroot-resistant hybrids on a clubroot-infested field may reduce the level of spores being released back into the soil from a canola crop, it will not control spores already contained in the soil. In other words, it will not prevent the spread from field to field through soil transfer. It's very important to remain diligent with sanitation practices, keeping footwear, tools, vehicles and equipment clean in order to slow the spread of clubroot. At a minimum, large chunks of soil should be removed from equipment prior to leaving an infested field. The best sanitation product is bleach in a 2% solution.

Q. Can clubroot be spread via seed?

A. Clubroot is not a seed-borne disease. Clubroot can only spread through resting spores contained in soil. That's why seed only plays a role in clubroot spread if earth tag is present on the seed surface. Earth tag are the soil particles that may be resting on the seed coat of harvested seed. The risk associated with disease spread via earth tag is very low. The primary means of disease spread is soil transfer on equipment—and represents a much greater risk. With a small-seeded crop such as canola, earth tag is not a significant issue and the cleaning and treating process further reduces risk. InVigor seed production takes a proactive approach to produce hybrid canola seed in fields not affected by clubroot. These steps include education of seed production staff, intensive field surveys prior to growing the seed crop, a minimum four-year rotation from brassica crops (canola, mustard and rapeseed) and a survey of every seed production field prior to harvest to confirm clubroot is not present.

Q. Are there any seed treatments or soil amendments available for clubroot?

A. Currently, there are no effective seed treatments or soil amendments available for controlling clubroot.

Q. Does rotating clubroot-resistant hybrids alone help prevent resistance?

A. Our recommendations on when to employ first- and second-generation genetics apply to all seed brands and not just InVigor. To date, there are limited sources of clubroot resistance available in commercial hybrids. Switching to a different canola brand does not ensure protection unless the resistance mechanisms are different and effective against the pathotypes that exist in your field. As more clubroot resistance sources are found, and a greater availability of genetic resistance diversity becomes available, then it makes sense to switch to a different source of resistance (or generation) that is effective against the pathotypes identified to cause the disease in your situation.

Q. Why doesn't BASF label their resistance mechanisms?

A. Currently, we label our clubroot-resistant hybrids as either containing first-generation and/or or second-generation resistance mechanisms. The science and understanding of this pathogen continues to evolve, and we're only starting to learn about clubroot population dynamics. Clubroot pathotypes within any given field or gall are more diverse than originally thought and current tests only generalize the predominant pathotype present. Due to this uncertainty and lack of specificity in pathogen identity, there is too great a risk in assigning genetic resistance to a particular field. In addition, since there are currently no commercial testing facilities that provide pathotyping for clubroot, understanding the specific details of clubroot in your field isn't a reality for most growers, so a more detailed description of the genetic resistance mechanisms present in a hybrid at this time will only add more confusion to an already complex problem.

Q. Are certain areas of the United States more prone to clubroot development?

A. Although clubroot is established primarily in the Langdon, North Dakota area, geography does not equate to immunity. Diligent scouting is recommended in all canola growing regions.

InVigor[®]

BASF
We create chemistry

Always read and follow label directions.

InVigor, Liberty and LibertyLink are registered trademarks of BASF. © 2024 BASF Corporation. All rights reserved.

ROUNDUP, ROUNDUP READY and TRUFLEX are registered trademarks of Bayer Group or its affiliated companies. Used under license. © 2024 Bayer Group. All rights reserved.

Pioneer brand canola products are registered trademarks of Corteva Agriscience and its affiliated companies.

All other trademarks are the property of their respective owners.