

Priaxor®

Xemium® Brand Fungicide

Use Guide for Cotton

Benefits of Priaxor® Fungicide

- Effective control of key foliar diseases
- Improves yield and minimizes risk
- Provides Plant Health benefits for cotton

Foliar Diseases are Found in Most Georgia Cotton Fields



Target Spot



Ascochyta Leaf Spot



Stemphylium Leaf Spot

Priaxor Fungicide Controls Foliar Diseases in Cotton



Untreated



Priaxor Fungicide

2014 – University of Georgia fungicide trial. Arlington, GA. Priaxor fungicide 4 fl oz/A.

Priaxor fungicide helps growers get the most out of every acre.



Technical Information Bulletin

150 years

 **BASF**
We create chemistry

Factors that Improve the Odds of Success and Yield Increases with Priaxor® Fungicide on Cotton

Foliar diseases of cotton are found in most cotton fields in Georgia and Alabama. Target Spot is especially widespread, can cause significant yield loss and is not linked to fertility deficiencies. Recent university field trials have shown Priaxor fungicide can reduce the severity of this disease and others, protect cotton leaves and improve yield.

If any of these factors below are present on your farm, applications of fungicides may be beneficial

- Have you or your neighbors had a problem within the past several years with disease on your cotton leaves? Symptoms include leaf spots and premature defoliation.
- Are you planting cotton behind cotton? Or on a short rotation?
- Is the weather favorable for disease development? Are afternoon showers in the forecast? Is a tropical storm forecast to impact the area?
- Is your cotton irrigated?
- Is your cotton managed for high yields?
- Is your cotton a full season variety?



Best Use Recommendations


- Use Rate: 4.0 fl oz/A
- Application Timing: 1st week of bloom; again 2–3 weeks later

Target Diseases

- Alternaria leaf spot and boll rot
- Anthracnose boll rot
- Ascochyta blight and boll rot
- Cercospora blight and leaf spot
- Target spot
- Diplodia boll rot
- Fusarium hard lock
- Phoma blight
- Rust
- Stemphylium leaf spot


Priaxor Fungicide Disease Control & Yield Benefits

% Defoliation from Target Spot



2014 LSU fungicide trial. Bloom application. Ratings taken 30 days after treatment.

Higher Yields with Priaxor Fungicide



2012 – 2014 Average of 8 replicated field trials. Auburn University, University of AR, University of GA, University of TN, LSU and BASF TX.