

Priaxor[®]
Xemium[®] Brand Fungicide

Headline AMP[®]
Fungicide

Grow Smart[™]
with **BASF**

Maximize Corn Yield Potential with BASF Fungicides for Disease Control and Plant Health

Benefits of Priaxor[®] Fungicide Applied Pre-Tassel

- Controls Early Season Diseases
- Increases Root Growth, Photosynthesis and Leaf Health
- Improves Stress Tolerance

Benefits of Headline AMP[®] Fungicide Applied at Tassel


- Delivers the Most Yield at Tassel
- Best in Class Preventative and Post-Infection Disease Control
- Improves Stalk Strength

On-Farm Multi-Hybrid x Fungicide Trial, Polo, IL 2015



↑ Headline AMP Fungicide at Tassel (VT) ↑ Priaxor Fungicide @ V5 fb Headline AMP Fungicide ↑ Priaxor Fungicide @ V5 ↑ No Fungicide Treatment

Dyna-Gro[®] Seed hybrids. Hybrids were replicated twice. Aerial photo taken 9/22/15.



$$\text{Yield} = \text{Ears/Acre} \times \text{Kernels/Ear} \times \text{Weight/Kernel}$$

Technical Information Bulletin

BASF
We create chemistry

Yield Results of Multiple Dyna-Gro® Hybrids to Various BASF Fungicides and Application Timings, Polo, IL 2015

	Yield (bu/A)			
	No Fungicide Treatment	Priaxor® Fungicide V5	Headline AMP® Fungicide VT	Priaxor Fungicide V5 fb Headline AMP Fungicide VT
All Dyna-Gro Seed Hybrids (n=6)	185	201	216	221
Dyna-Gro D40SS48 (100 RM)	183	185	196	204
Dyna-Gro D46SS62RIB (106 RM)	201	215	235	231
Dyna-Gro D47SS23RIB (107 RM)	191	197	211	214
Dyna-Gro D48SS76 (108 RM)	164	193	208	214
Dyna-Gro D50SS43 (111 RM)	176	199	212	221
Dyna-Gro D52SS91 (112 RM)	192	214	236	243

Best Use Recommendations for Priaxor Fungicide

- **Use Rate:** 4 fl oz/A
- **Labeled Crops:** Corn (all types)

Application Information

- Aerial: 2 GPA minimum; Ground: 10 GPA minimum
- PHI: 7 days for sweet corn; 21 days for all other types
- REI: 12 hours

Adjuvants

- Adjuvant flexible; however, see label for adjuvant restrictions after the V8 stage and prior to the VT stage of corn growth

Target Diseases

- Anthracnose
- Eyespot
- Gray leaf spot
- Northern corn leaf blight
- Northern corn leaf spot
- Physoderma brown spot
- Rust, southern and common
- Southern corn leaf blight
- Yellow leaf blight

Best Use Recommendations for Headline AMP Fungicide

- **Use Rate:** 10 to 14.4 fl oz/A
- **Labeled Crops:** Corn (all types)

Application Information

- Aerial: 2 GPA minimum; Ground: 10 GPA minimum
- PHI: 20 days for field and popcorn grain; 7 days for seed and sweet corn
- REI: 12 hours

Adjuvants

- Adjuvant flexible; however, see label for adjuvant restrictions after the V8 stage and prior to the VT stage of corn growth

Target Diseases

- Anthracnose
- Eyespot
- Gray leaf spot
- Northern corn leaf blight
- Northern corn leaf spot
- Physoderma brown spot
- Rust, southern and common
- Southern corn leaf blight
- Yellow leaf blight

Priaxor®
Xemium® Brand Fungicide

Headline AMP®
Fungicide



Always read and follow label directions.

Grow Smart is a trademark and Priaxor and Headline AMP are registered trademarks of BASF. Dyna-Gro is a registered trademark of Crop Production Services.

©2015 BASF Corporation. All Rights Reserved. APN# 1512002 Headline AMP-Priaxor-Corn-CPS Hybrids

BASF
We create chemistry