

Priaxor®

Xemium® Brand Fungicide

Use In California Tomatoes

Benefits of Priaxor® Fungicide

- Driven by Xemium® Fungicide technology advantages
 - Broader activity across disease lifecycles
 - Continuous redistribution and extended residual protection
- Contains F500® – the #1 Plant Health Fungicide
- More consistent disease control helps maximize yield and quality potential

Priaxor provides more consistent disease control to help maximize yield and quality potential.

Priaxor Fungicide Use in Tomatoes

Use Rates and Diseases	4–8 fl oz/A: Black mold, Early blight, Septoria leaf spot, Target spot and Anthracnose
	6–8 fl oz/A: Powdery mildew
	8 fl oz/A: Suppression Botrytis gray mold, Sclerotinia stem rot, White mold, and Late blight
PHI	0 days
REI	12 hours



150 years

 **BASF**

We create chemistry

Technical Information Bulletin

Best Management Practices

Adjuvant Use and Tank Mixing with Priaxor® Fungicide

- Always consult a BASF representative before tank mixing products with Priaxor fungicide
- Effectiveness and safety of Priaxor fungicide in mixes depends on product, growth stage and other factors
- Do not mix with EC, COC, MSO, OS, MSO/OS blends
- NIS can be used at their lowest labeled rates up to 0.125%



Black Mold



Powdery Mildew



Early Blight




Anthracnose




Disease Control and Plant Productivity

Powdery Mildew Control



2014 Brenna Aegerter – UCCE, San Joaquin County. Rating 9/18/14. Applications made 8/22 & 9/5. 44 GPA in CO2 sprayer at 40 psi. NIS @ 0.25% with Priaxor fungicide, NIS 0.125% v/v with Quadris Top. Fresh Market Variety: Valley Cat.

Tomato Yield and Quality



2014 Brenna Aegerter – UCCE, San Joaquin County. Applications made 8/22, 9/5, & 9/19. 44 GPA in CO2 sprayer at 40 psi. NIS @ 0.25% with Priaxor fungicide, NIS 0.125% v/v with Quadris Top. Fresh Market Variety: Valley Cat.