# Beyond<sup>®</sup> Xtra

Herbicide

## Alfalfa

## **Best Use Recommendations**

#### Crop\*

## **Recommended Timing**

### Alfalfa

**Herbicide timing\*:** Apply early post-emergence when weeds are actively growing and before they exceed a height of 3 inches.

- Seedling alfalfa: Apply when seedling alfalfa is in the second trifoliate stage or larger and when the majority of weeds are 1-inch to 3-inches tall. When applied to alfalfa grown for seed, apply before bud formation.
- **Established alfalfa:** Apply to established alfalfa in fall, winter, or spring to dormant or semidormant alfalfa, or between cuttings. Apply before significant alfalfa growth or regrowth (3-inches).

## **Active Ingredients:**

Imazamox - Group 2

#### Formulation:

Soluble liquid (SL)

#### **Product Packaging:**

2 x 1 gallon

## Storage:

Do not store below 32°F

#### **Bulk Density:**

8.75 lbs/gal

Crop	Weeds Controlled <sup>†</sup>	Application Rate
Alfalfa	Broadleaf weeds  Lambsquarters, common  Pigweed, redroot  Nightshade spp.  Mustard spp. (black, tumble, wild)  Thistle, Russian  Buckwheat, wild‡  Kochia‡	4-6 fl oz/A
	Grass weeds ■ Foxtail spp. (green, yellow) ■ Oat, wild <sup>‡</sup>	

- \* See label for complete listing of labeled crops.
- † See label for complete listing of weeds controlled.
- ‡ Beyond Xtra herbicide does not control ALS-resistant weeds.



#### **Application Tips**

Rainfast: 1 hour REI: 4 hours PHI: 0 days\*

Nozzle, droplet size

- Flat fan
- Ground app minimum of 10 GPA; 20-40 PSI
- Ground app w/ low volume sprayer minimum of 10 GPA; 40-60 PSI
- Aerial app minimum 5 GPA

#### Seasonal max use rate: 6 fl oz/A

\*Application recommendations vary by region and crop. Please contact your local BASF representative and/or refer to the product label for additional details.

#### **Mixing Order**

- 1. **Water:** Begin by agitating a thoroughly clean sprayer tank ½ ¾ full of clean water
- 2. **Agitation:** Maintain constant agitation throughout mixing and application
- 3. Soluble-packet products and thoroughly mix
- 4. **WP, DG, DF, or liquid flowable formulations** not in soluble packets
- 5. Beyond® Xtra herbicide and thoroughly mix
- 6. Other aqueous solution products
- 7. EC products
- 8. Surfactant or crop oil
- 9. Nitrogen fertilizer solution
- 10. Remaining quantity of water; maintain agitation

#### **Adjuvants, Fertilizers and Tank Mixes**

Crop	Rate	Adjuvants	Fertilizers	Additional Comments <sup>†</sup>
Newly Seeded Alfalfa	5 fl oz/A	NIS @ 1% v/v	AMS @ 12-15 lbs/100 gal	For best results, use at least 20 GPA for optimal coverage.
Established Alfalfa	5-6 fl oz/A	COC or MSO @ 1% v/v	AMS @ 12-15 lbs/100 gal	Apply before alfalfa has 3" of re-growth to insure adequate coverage on weeds
Tough Grasses in Newly Seeded or Established Alfalfa	5 fl oz/A	COC @ 1% v/v	AMS @ 12-15 lbs/100 gal	Apply before alfalfa has 3" of re-growth or to newly seeded alfalfa with at least two fully expanded trifoliates to insure adequate coverage.

<sup>†</sup> DO NOT tank mix organophosphate or carbamate insecticides with Beyond Xtra herbicide on listed crops unless otherwise specified in writing by BASF. Refer to product label for more use restrictions.







