

Beyond[®] Xtra

Herbicide

Clearfield[®] Rice

Best Use Recommendations

Crops*

Recommended Timing

Clearfield rice

Herbicide timing: Apply early post-emergence when weeds are actively growing and before broadleaf weeds exceed a height of 3 inches and grass weeds exceed 4-5 leaves.

- **Clearfield rice varieties:** 4-leaf to rice panicle initiation (green ring) plus 14 days.
- **Clearfield rice hybrids:** 4-leaf to rice panicle initiation. **DO NOT** apply Beyond[®] Xtra herbicide to Clearfield rice hybrids after panicle initiation.

Active Ingredients:

Imazamox – Group 2

Formulation:

Soluble liquid (SL)

Product Packaging:

2 x 1 gallon

Storage:

Do not store below 32°F

Bulk Density:

8.75 lbs/gal

Crops	Weeds Controlled [†]	Application Rate
Clearfield rice	<p>Broadleaf weeds</p> <ul style="list-style-type: none">■ Morning glory spp. (entireleaf, ivyleaf, smallflower, tall)■ Smartweed spp. (ladysthumb, Pennsylvania, swamp) <p>Grass weeds</p> <ul style="list-style-type: none">■ Barnyardgrass■ Sprangletop■ Signalgrass, broadleaf■ Panicum, fall■ Crabgrass, large■ Rice, red[‡]	4-6 fl oz/A**

*See label for complete listing of labeled crops.

**Beyond Xtra herbicide can only be applied following at least one application of Newpath[®] herbicide (imazethapyr) or Clearpath[®] herbicide (imazethapyr and quinclorac). DO NOT make more than two applications of Beyond Xtra herbicide per year (growing season).

† See label for complete listing of weeds controlled.

‡ See label for specific application rate & timing details.

 **BASF**

We create chemistry

Application Tips

Rainfast: 1 hour

REI: 4 hours

PHI: 0 days

Adjuvant**

- 1-2 pts/A of COC **REQUIRED**

Nozzle, droplet size

- Flat fan
- Ground app – minimum of 10 gpa; 20-40 psi
- Ground app w/ low volume sprayer – minimum of 10 gpa; 40-60 psi
- Aerial app – minimum 5 gpa

Seasonal max use rate: 10 fl oz/A

**Application recommendations vary by region and crop. Please contact your local BASF representative and/or refer to the product label for additional details.

Tank Mixes and Additives***

DO NOT tank mix organophosphate or carbamate insecticides with Beyond® Xtra herbicide on listed crops unless otherwise specified in writing by BASF. For tank mixing, refer to respective product labels for restrictions.

***Refer to product label for more use restrictions.

 To learn more about crop protection products from BASF, visit www.agproducts.basf.us

Mixing Order

1. **Water:** Begin by agitating a thoroughly clean sprayer tank $\frac{1}{2}$ – $\frac{3}{4}$ full of clean water
2. **Agitation:** Maintain constant agitation throughout mixing and application
3. **Soluble-packet products** and thoroughly mix
4. **WP, DG, DF, or liquid flowable formulations** not in soluble packets
5. **Beyond® Xtra herbicide** and thoroughly mix
6. **Other aqueous solution products**
7. **EC products**
8. **Surfactant or crop oil**
9. **Nitrogen fertilizer solution**
10. **Remaining quantity of water;** maintain agitation



 **BASF**

We create chemistry

Beyond® Xtra
Herbicide