

Beyond[®] Xtra

Herbicide

Clearfield[®] and Clearfield Plus Sunflower

Best Use Recommendations

Crop*

Recommended Timing

Clearfield/Clearfield Plus sunflower

Herbicide timing: Early postemergence (2-leaf to 8-leaf sunflower stage); apply before grass weeds exceed 4-5 leaves and broadleaf weeds exceed a height of 3 inches.

Active Ingredients:

Imazamox – Group 2

Formulation:

Soluble liquid (SL)

Product Packaging:

2 x 1 gallon

Storage:

Do not store below 32°F

Bulk Density:

8.75 lbs/gal

Crop	Weeds Controlled†	Application Rate
Clearfield/Clearfield Plus sunflower	<p>Broadleaf weeds</p> <ul style="list-style-type: none">■ Lambsquarters, common■ Pigweed, redroot■ Nightshade spp. (black, Eastern black, hairy)■ Mustard, wild■ Buckwheat, wild <p>Grass weeds</p> <ul style="list-style-type: none">■ Foxtail spp. (giant, green, yellow)■ Barnyardgrass■ Oat, wild‡■ Brome, downy‡■ Jointed goatgrass‡	<p>Clearfield sunflower 4 fl oz/A</p> <p>Clearfield Plus sunflower 4-6 fl oz/A</p>

* See label for complete listing of labeled crops.

† See label for complete listing of weeds controlled.

‡ Beyond Xtra herbicide does not control ALS-resistant weeds.



BASF

We create chemistry

Application Tips

Rainfast: 1 hour

REI: 4 hours

PHI: 0 days

Adjuvants, Fertilizers and Tank Mixes**

Crop	Adjuvants	Fertilizers
Clearfield® sunflower	Include NIS or Basic Blend Adjuvant; DO NOT use COC, MSO, OR HSOC	Apply with AMS or UAN
Clearfield Plus sunflower	Include NIS, COC, MSO, or HSOC†	Apply with AMS or UAN

† Use of COC, MSO, or HSOC in place of NIS in Clearfield Plus sunflower may increase crop response.

Nozzle, droplet size

- Flat fan
- Ground app – minimum of 10 GPA; 20-40 PSI
- Ground app w/ low volume sprayer – minimum of 10 GPA; 40-60 PSI
- Aerial app – minimum 5 GPA

**Application recommendations vary by region and crop. Please contact your local BASF representative and/or refer to the product label for additional details.



To learn more about crop protection products from BASF, visit www.agproducts.basf.us

Tank Mixes and Additives***

DO NOT tank mix organophosphate or carbamate insecticides with Beyond® Xtra herbicide on listed crops unless otherwise specified in writing by BASF.

***Refer to product label for more use restrictions.

Mixing Order

1. **Water:** Begin by agitating a thoroughly clean sprayer tank ½ – ¾ full of clean water
2. **Agitation:** Maintain constant agitation throughout mixing and application
3. **Soluble-packet products** and thoroughly mix
4. **WP, DG, DF, or liquid flowable formulations** not in soluble packets
5. **Beyond Xtra herbicide** and thoroughly mix
6. **Other aqueous solution products**
7. **EC products**
8. **Surfactant or crop oil**
9. **Nitrogen fertilizer solution**
10. **Remaining quantity of water;** maintain agitation



BASF
We create chemistry

Beyond® Xtra
Herbicide