

# Facet® L

Herbicide


## Strong and Flexible: For Grass and Broadleaf Weed Control in Rice

### Benefits of Facet® L Herbicide

- Convenient, efficacious liquid formulation
- Consistent control of resistant annual grasses and broad leaf weeds including: hemp sesbania, joint vetches, and propanil-resistant barnyardgrass

### Greater Consistency

Maintains Excellent Preemergence Control of Barnyardgrass



BASF Field Trial (2009): Winchester, AR



**Facet L herbicide provides consistent control of resistant annual grasses and broadleaf weeds.**


### General Information

#### Labeled Crop: Rice

- Can be applied to the soil surface before, during or after planting dry-seeded rice.
- When applied as a foliar spray, rice must be in at least the 2-leaf stage. For best control, establish permanent flood within 2 days after Facet L herbicide application.

### Faster Acting

Simulated rainfall 4 hours after application



BASF Greenhouse Trial (2005), RTP, NC

**BASF**

We create chemistry

**Technical Information Bulletin**


# Facet® L Herbicide Activity on Key Weeds in Rice

Weed Species	Soil Applications (Rate per Acre)			Foliar Applications (Rate per Acre)	
	Light-textured: sandy loams	Medium-textured: silts, loams, silt loams, sandy clay loams	Heavy-textured: silty clays, silty clay loams, clay loams, clays, gumbo and buckshot	In postemergence applications of Facet L herbicide, add 2 pints of crop oil concentrate per acre for improved uptake and enhanced weed control.	
				Small weeds controlled and short-term soil residual	Larger weeds controlled and long-term soil residual
<b>Annual Grasses</b> Barnyardgrass Broadleaf signalgrass Junglerice Large Crabgrass	21 to 28 fl ozs	32 fl ozs	42 fl ozs	25 to 32 fl ozs up to 2 inches	25 to 43 fl ozs up to 3 inches
<b>Broadleaf Weeds</b> Eclipta Jointvetch species Indian Northern Morningglory species cypressvine entireleaf ivyleaf palmleaf pitted purple moonflower tall (common) Sesbania, hemp	21 to 28 fl ozs	32 fl ozs	42 fl ozs	25 to 32 fl ozs up to 2 leaves	32 to 42 fl ozs up to 3 leaves
Alligatorweed (partial control)*	n/a	n/a	n/a	43 fl ozs	n/a

\*When applied as a foliar spray, rice must be in at least the 2-leaf stage. For best control, establish permanent flood within 2 days after Facet L herbicide application.

## Innovative Formulation

- Excellent handling characteristics
- Easy mixing
- Excellent tankmix compatability




Quinclorac DF Formulation      Facet L Herbicide

## Best Use Recommendations

Comparable Rates	
Quinclorac DF Formulation lbs per Acre	Facet 1.5 L Herbicide fl ozs per Acre
0.33	21
0.40	26
0.44	28
0.50	32
0.67	43

In postemergence applications of Facet L and Quinclorac DF Formulation, add 2 pints of crop oil concentrate per acre for improved uptake and enhanced weed control.

## Flexibility, Wide Application Window



**BASF**  
We create chemistry

**Facet® L**  
Herbicide